

Background on the Minnesota Tobacco Withdrawal Scale-Revised (MTWS-R)
(<http://www.med.uvm.edu/behaviorandhealth/minnwsdefault>)

John R. Hughes, M.D.
(john.hughes@uvm.edu)

2/26/20

1. I have changed the name of the scale to the Minnesota Tobacco Withdrawal Scale to reflect that non-nicotine factors can influence scores. Several versions of the MTWS have been used with different numbers of items and response options. The enclosed scales are the versions I recommend.
2. Six articles have reviewed the psychometrics of the scale ¹⁻⁷. Five articles have comprehensively reviewed what is known about tobacco withdrawal^{1, 2, 4, 8, 9}. Other articles have pointed out optimal methods for measuring withdrawal ^{4, 10, 11}.
3. The seven other commonly used withdrawal scales are the Cigarette Withdrawal Scale¹² (<https://eprovide.mapi-trust.org/instruments/cigarette-withdrawal-scale>), the Mood & Physical Symptoms Scale¹³ (www.robertwest@ucl.ac.uk), subscales of the Profile of Mood States Manual¹⁴ (www.mhs.com), the Shiffman Jarvik Withdrawal Scale¹⁵(shiffman@pinneyassociates.com), the Smoker Complaints Scale¹⁶ (ngs@ucla.edu), the Wisconsin Smoking Withdrawal Scale¹⁷ (<https://eprovide.mapi-trust.org/instruments/wisconsin-smoking-withdrawal-scale>) and a scale to use on a hand-held computer¹⁸ (shiffman@pinneyassociates.com). Three studies have empirically compared our scale with those of others⁵⁻⁷
4. Attached is a table comparing our scale with the criteria for Nicotine/Tobacco Withdrawal from the American Psychiatric Association's proposed Diagnostic and Statistical Manual, Fifth Edition-Revised (DSM-V)¹⁹ and the World Health Organization's International Classification of Disease, Tenth Edition (ICD-10)²⁰ and other withdrawal scales.
5. The scale is not copyrighted and you do not need our permission to use it.
6. There are no true population norms. Estimates for the prevalence of withdrawal for US, Germany and Asia populations have been described^{8, 21}

Reference List

- (1) Piper ME. Withdrawal: Expanding a key addiction construct. *Nicotine & Tobacco Research* 2015; 17:1405-1415
- (2) Piper ME, McCarthy DE, Baker TB. Assessing tobacco dependence: A guide to measure evaluation and selection. *Nicotine and Tobacco Research* 2006;8:339-351.
- (3) Shiffman S, West RJ, Gilbert DG. Recommendation for the assessment of tobacco craving and withdrawal in smoking cessation trials. *Nicotine & Tobacco Research* 2004;6:599-614.
- (4) Hughes JR. Measurement of the effects of abstinence from tobacco. A qualitative review. *Psychol Addict Behav* 2007;21:127-137.
- (5) West R, Ussher M, Evans M, Rashid M. Assessing DSM-IV nicotine withdrawal symptoms: A

comparison and evaluation of five different scales. *Psychopharm* 2006;184:619-627.

- (6) Etter JF, Hughes JR. A comparison of the psychometric properties of three cigarette withdrawal scales. *Addiction* 2006;101:362-372.
- (7) Javitz H, Lerman C, Swan G. Comparative dynamics of four smoking withdrawal symptom scales. *Addiction* 2012;107:1501-1511.
- (8) Hughes JR. The effects of abstinence from tobacco: Etiology, determinants, epidemiology, significance. *Nicotine & Tobacco Research* 2007;9:329-339.
- (9) Hughes JR. The effects of abstinence from tobacco: Valid symptoms and time course. *Nicotine & Tobacco Research* 2007;9:315-327.
- (10) Hughes JR. Distinguishing withdrawal relief and direct effects of smoking. *Psychopharm* 1991;104:409-410.
- (11) Hughes JR, Hatsukami DK. Errors in using tobacco withdrawal scales. *Tobacco Control* 1998;7:92-93.
- (12) Etter J. Self-administered questionnaire to measure cigarette withdrawal symptoms: The Cigarette Withdrawal Scale. *Nicotine & Tobacco Research* 2005;7:47-57.
- (13) West R, Hajek P. Evaluation of the Mood and Physical Symptoms Scale (MPSS) to assess cigarette withdrawal. *Psychopharm* 2004;177:195-199.
- (14) McNair DM, Lorr J, Droppelman LF. *Profile of Mood States Manual*. North Tonawanda, NY: Multi-Health Systems; 1992.
- (15) Shiffman S, Jarvik ME. Smoking withdrawal symptoms in two weeks of abstinence. *Psychopharmacologia* 1976;50:35-39.
- (16) Schneider NG, Jarvik ME. Time course of smoking withdrawal symptoms as a function of nicotine replacement. *Psychopharm* 1984;82:143-144.
- (17) Welsch SK, Smith SS, Wetter DW, Jorenby DE, Fiore MC, Baker TB. Development and validation of the Wisconsin Smoking Withdrawal Scale. *Exp Clin Psychopharm* 1999;7(4):354-361.
- (18) Shiffman S, Paty JA, Gnys M, Kassel JA, Hickcox M. First lapses to smoking: Within-subjects analysis of real-time reports. *J Consult Clin Psychol* 1996;64:366-379.
- (19) American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 5th ed. Washington: American Psychiatric Association; 2013.
- (20) World Health Organization. *The ICD-10 Classification of Mental and Behavioural Disorders. Diagnostic Criteria for Research*. Geneva, Switzerland: World Health Organization; 1993.
- (21) Hughes JR, Helzer JE, Lindberg SA. Prevalence of DSM/ICD-defined nicotine dependence. *Drug Alcohol Depend* 2006;85:91-102.