



LIVING & WORKING IN PRIMARY CARE DURING COVID-19

A research study to understand how the primary care professional is personally affected during this crisis

Results:


59%

of participants are experiencing burnout in one or more domains. Emotional exhaustion and diminished feelings of accomplishment are the most common experiences of burnout.

Nearly **half** of participants indicated that their work schedule did NOT leave them enough time for their personal/family life. Not having enough time for personal/family life was highly predictive of burnout.

Methods: REDCap online surveys were distributed to primary care professionals weekly starting May 18, 2020. Participants can choose to complete the survey once or several times.

Participants: As of April 12, 2021, we have recruited 1,153 participants from all 50 states: 37% physicians, 12% nurse practitioners and physician assistants, 15% nurses, 7% medical assistants, 13% behavioral health providers, 6% non-clinical, and 10% other clinical; 78% of participants are women, 86% white, and the average years working in their current role is 8 years (median; 0.1 to 70 years).



Contact Us: 802-656-4560 ° healthcare.covid19@med.uvm.edu ° go.uvm.edu/primarycare

Note: *Data cleaning has resulted in the removal of potential duplicates, incomplete responses, and ineligible participants (total of 43 removed from analysis to date). Interpret preliminary results with caution. The findings are not generalizable given the small sample size. In other words, we cannot assume these data reflect all primary care professionals. The number of respondents may change based on the questions posed for the week/month. The aMBI (Riley, et al., 2017) was modified to assess acute (weekly) burnout (range of each subscale: 0-12). Scoring: Exhaustion 6+; Depersonalization 3+; Accomplishment <7. Therefore, the results cannot be compared across studies using the aMBI. For tips/resources on coping with distress during a pandemic, click [here](#).



Note: *Data cleaning has resulted in the removal of potential duplicates, incomplete responses, and ineligible participants (total of 43 removed from analysis to date). Interpret preliminary results with caution. The findings are not generalizable given the small sample size. In other words, we cannot assume these data reflect all primary care professionals. The number of respondents may change based on the questions posed for the week/month. The aMBI (Riley, et al., 2017) was modified to assess acute (weekly) burnout (range of each subscale: 0-12). Scoring: Exhaustion 6+; Depersonalization 3+; Accomplishment <7. Therefore, the results cannot be compared across studies using the aMBI. For tips/resources on coping with distress during a pandemic, click [here](#).